

Fig 1. Mesomeric forms of p-benzosemiquinone. (a) anionic; (b),(c) neutral; (d) cationic.

10076033.021302




Fig 2. Band thin layer chromatography of the methanol solution after lyophilization (step 5). —> Indicates the band of cs-oxidant.




Fig 3. HPLC profile the butanol extract after TLC. The cs-oxidant (step 6) eluted as a major peak at the retention time of 8.808 min. The amount of cs-oxidant eluted was $\approx 12 \mu\text{g}$.

20251201 E034007




Fig 4. HPLC profile of the pure cs-oxidant, eluted at the retention time of 8.808 min.




Fig 5. Thin layer chromatography of the pure cs-oxidant ($R_f = 0.26$)

[illegible]

Fig 6a. Fluorescence spectroscopic profile of the cs-oxidant in methanol. The excitation was at 293 nm and the emission scanning was measured from 300 nm to 800 nm. The emission maxima were at 329.6 nm and at 651.4 nm.

PEAK	TIME (min)	AREA	HEIGHT	WIDTH
1	0.514	1.114	1.114	1.114
2	1.014	3.750	3.750	3.750
3	1.514	1.114	1.114	1.114
4	2.514	1.114	1.114	1.114
5	3.514	3.750	3.750	3.750

$\frac{1}{2} \times 100 = 50$

NO.	PEAK	WAVELENGTH	WAVELENGTH	WAVELENGTH
1	228.2nm	330 nm	248nm/min	77.40
2	293.8nm	2 sec		

Fig 7a. Fluorescence spectroscopic profile of the cs-oxidant in methanol. The emission was at 330 nm and the excitation scanning was measured from 220 nm to 325 nm. The excitation maxima were at 228.2 nm and at 293.8 nm.

[illegible]

Fig 7a. Fluorescence spectroscopic profile of the cs-oxidant in methanol. The emission was at 330 nm and the excitation scanning was measured from 220 nm to 325 nm. The excitation maxima were at 228.2 nm and at 293.8 nm.




Fig 7b. Fluorescence spectroscopic profile of the cs-oxidant in methanol. The emission was at 651 nm and the excitation scanning was measured from 220 nm to 650 nm. The excitation maxima were at 229.2 nm and at 294.8 nm.




Fig 8. Crystal structure of the pure cs-oxidant

2026033 02302




Fig 9. UV spectrophotometric profile of the cs-oxidant in methanol. It has two absorption maxima, one at 293.4 nm and another at 223.0 nm.




Fig.10 Stability of the solid oxidant kept at 25°C under darkness. The stability was determined by its capacity to oxidize ascorbic acid. Ascorbic acid was measured by HPLC analysis at 254 nm.




Fig.11 Standard curve of ascorbic acid based on HPLC analysis at 254 nm.

20075033-034302




Fig.12 Stability of the cs-oxidant in 50 mM potassium phosphate buffer at 25°C measured by its potency to oxidize ascorbate as evidenced by HPLC area.




Fig.13 Quantitative reduction of ferricytochrome c by the oxidant as measured by the formation of ferrocytochrome c with time at 550 nm. The reaction was carried out in 50 mM potassium phosphate buffer, pH 7.4, keeping the final concentration of ferricytochrome c at 100 μ M. One nmole of the oxidant reduced 0.71 nmole of ferricytochrome c.

Amount of the ox-oxidant (μg)	Absorbance at 550 nm
1	0.16
2	0.32
3	0.49
4	0.65
5	0.82

Fig 15. Standard curve of the oxidant on the basis of reduction of cytochrome c by using different amounts of the oxidant ranging from 1 μ g to 5 μ g.






Fig 16. Mass spectrum of the pure cs-oxidant

Fig 17. UV spectrophotometric profile of the hydroquinone in methanol. It has two absorption maxima, one at 293.8 nm and another at 224.2 nm.




10076033 024302



SPEED: 120.0 nm/min RESPONSE: MEDIUM
 PASS: 2.00nm
 NO. PEAK VALLEY
 1 293.6 nm 0.2772 Abs 252.8 nm 0.0269 Abs
 2 224.4 nm 0.5476 Abs 214.0 nm 0.4314 Abs

Fig 18. UV spectrophotometric profile of the cs-oxidant stored at room temperature in dark for 8 days. The two absorption maxima are at 293.6 nm and at 224.4 nm.

2024.03.20 14:00:00



SCAN SPEED: 120.0 nm/min RESPONSE: MEDIUM
 WAVELENGTH: 2.00nm

NO.	PEAK		VALLEY	
	nm	Abs	nm	Abs
1	263.4	0.1407	209.6	0.8263

Fig 19. UV spectrophotometric profile of equimolar mixture of p-benzoquinone and hydroquinone in methanol. There is a shoulder near 242 nm (the λ_{max} of p-benzoquinone).




Fig 20. Fluorescence spectroscopic profile of the hydroquinone in methanol. The excitation was at 294 nm and the emission scanning was measured from 300 nm to 800 nm. The emission maxima were at 329.4 nm and at 651.6 nm.

202120-0009-0001




Fig 21. FTIR spectroscopic profile of the cs-oxidant.

20251201 1007503.043001




Fig 22 FTIR spectroscopic profile of hydroquinone




Fig 23 H-NMR spectroscopic profile of the cs-oxidant in CD_3COCD_3

20070303 0400




Fig 24. H-NMR spectroscopic profile of hydroquinone in CD_3COCD_3

10076033.024302
202120 EE09/00T




Fig 25 Comparative H-NMR spectroscopic profiles of (a) cs-oxidant and (b) hydroquinone.

20E120" E09200T




Fig 26. H-NMR spectroscopic profile of the cs-oxidant after reduction with sodium dithionite.

20250701 10:00




Fig 27 ¹³C-NMR spectroscopic profile of the cs-oxidant in CD₃COCD₃

00120 66094001




Fig 28. C-NMR spectroscopic profile of hydroquinone in CD₃COCD₃




Fig 29. Room temperature ESR spectrum of CS-oxidant, freshly prepared from 100 cigarettes. The spectrum was recorded on a JES-REIX ESR spectrometer (Tokyo, Japan). The spectral parameters were as follows: microwave frequency, 9.435 GHz; power, 2mW; field modulation width, 0.4mT; modulated frequency, 100kHz; time constant, 0.3 sec; scan rate, 2.5 mT/ sec

201201-021302




Fig 30. Room temperature ESR spectrum of aged (10 days) cs-oxidant, prepared from 400 cigarettes




Fig 31. HPLC profile of the whole cs solution analyzed in the silica column (Lichrospher® Si 60, Merck).

* Indicates the retention time, area and the concentration (13.6682%) of the cs-oxidant




Fig 32. HPLC profile of the aqueous extract of cs solution analyzed in the silica column (Lichrospher® Si 60, Merck).

* Indicates the retention time, area and the concentration (27.2269%) of the cs-oxidant




Fig 33. HPLC profile of the whole cs solution analyzed in the ODS column (Shim-pack CLC-ODS, Shimadzu). The cs-oxidant eluted at 13.467 min.

20070707 000000




Fig 34. HPLC profile of the pure cs-oxidant, analyzed in the ODS column (Shim-pack CLC-ODS, Shimadzu) eluted at the retention time of 13.458 min.




Fig 35a. SDS-PAGE of the guinea pig lung microsomal proteins treated with whole cs solution and the cs-oxidant. Lane 1, untreated microsomes; lane 2, microsomes treated with 50 μ l cs solution; lane 3, microsomes treated with 100 μ l cs solution; lane 4, microsomes treated with 10 μ g cs-oxidant; lane 5, microsomes treated with 20 μ g cs-oxidant.




Fig 35b. Densitometric scanning of the protein bands of different lanes as in Fig 35a.